

NT@UW-99-49

## Light quark distributions in the proton sea

Mary Alberg<sup>a\*</sup> and Ernest M. Henley<sup>b</sup><sup>a</sup>Department of Physics, Seattle University, Seattle, WA 98122, USA and  
Department of Physics, University of Washington, Seattle, WA 98195, USA<sup>b</sup>Department of Physics, University of Washington, Seattle, WA 98195, USA and  
Institute for Nuclear Theory, University of Washington, Seattle, WA 98195, USA

We use the meson cloud model to calculate  $\bar{d}(x) - \bar{u}(x)$  and  $\bar{d}(x)/\bar{u}(x)$  in the proton. We show that a modification of the symmetric, perturbative part of the light quark sea provides better agreement with the ratio  $\bar{d}(x)/\bar{u}(x)$ .

**1. Introduction**

Flavor asymmetry in the light quark sea of the proton has been well-established by experiment [1–4]. The violation of the Gottfried sum rule found by NMC [1] indicated that  $D \equiv \int_0^1 dx (\bar{d}(x) - \bar{u}(x)) = 0.148 \pm 0.039$ . In Drell-Yan experiments,  $\bar{d}/\bar{u}$  was determined to be greater than 2 at  $x = .18$  by NA51 [2], and the  $x$ -dependence of this ratio has recently been measured by E866 [3] in the region  $0.02 \leq x \leq 0.345$ . From their data and global parton distributions [5] E866 also determined the difference  $\bar{d}(x) - \bar{u}(x)$  and  $D = 0.100 \pm 0.018$ . Flavor asymmetry consistent with these results has also been seen in the measurement of  $\bar{d}(x) - \bar{u}(x)$  by HERMES [4].

It was first suggested by Thomas [6], and later by Henley and Miller [7] that a natural explanation for this asymmetry is the meson cloud of the proton. The net positive charge of the cloud leads to an excess of  $\bar{d}$  over  $\bar{u}$ . Other causes for the asymmetry have been invoked, such as antisymmetrization [8,9], but these have been insufficient to describe the data. For reviews, see Refs. [10,11].

We wish to emphasize that the E866 measurements provide a critical test of our understanding of the flavor-symmetric (FS) contributions to the light quark sea, as well as the flavor-asymmetric (FA) contributions. Meson cloud models [9,12–15] provide a reasonably good description of the  $\bar{d}(x) - \bar{u}(x)$  asymmetry, which depends on FA contributions alone, but fail to explain the broad maximum in the ratio  $\bar{d}(x)/\bar{u}(x)$  at  $x \approx 0.18$  and its return to unity, or even lower values, at larger  $x$ . FS terms also contribute to this ratio

$$\frac{\bar{d}(x)}{\bar{u}(x)} = \frac{\bar{d}(x) - \bar{u}(x)}{\bar{u}(x)} + 1, \quad (1)$$

---

\*Contributions to the work described in this talk have been made by students Thomas Falter (University of Giessen) and Adam Graunke (Seattle University).

and it is clear that  $\bar{u}(x)$  falls too rapidly with  $x$  in these models. We have proposed [16] that agreement with data can be improved by using harder distributions for the perturbative contributions to  $\bar{u}(x)$ , motivated by their origin in gluon splitting.

## 2. Pion cloud model

To illustrate this argument we use a pion cloud model. Other states should be included in a full calculation, but e.g. the  $\rho$  meson increases  $\bar{d}(x) - \bar{u}(x)$ , whereas the intermediate  $\Delta$  decreases it, so these effects tend to cancel. The wave function of the proton is written in terms of a Fock state expansion

$$|p\rangle = \sqrt{Z} |p\rangle_{\text{bare}} + \sum_{MB} \int dy d^2\vec{k}_\perp \phi_{BM}(y, k_\perp^2) |B(y, \vec{k}_\perp) M(1-y, -\vec{k}_\perp)\rangle, \quad (2)$$

with  $BM = p\pi^0, n\pi^+$ . The factor  $\sqrt{Z}$  is a wavefunction renormalization constant and  $\phi_{BM}(y, k_\perp^2)$  is the probability amplitude for finding a physical nucleon in a state consisting of a baryon  $B$  with longitudinal momentum fraction  $y$ , transverse momentum  $\vec{k}_\perp$ , and a meson  $M$  of momentum fraction  $(1-y)$ , transverse momentum  $-\vec{k}_\perp$ . The quark distribution functions  $q(x)$  in the proton are given by

$$q(x) = q^{\text{bare}}(x) + \delta q(x), \quad (3)$$

with

$$\delta q(x) = \sum_{MB} \left( \int_x^1 f_{MB}(y) q_M\left(\frac{x}{y}\right) \frac{dy}{y} + \int_x^1 f_{BM}(y) q_B\left(\frac{x}{y}\right) \frac{dy}{y} \right), \quad (4)$$

$$f_{BM}(y) = \int_0^\infty |\phi_{BM}(y, k_\perp^2)|^2 d^2k_\perp, \quad (5)$$

and

$$f_{MB}(y) = f_{BM}(1-y). \quad (6)$$

The splitting function  $f_{n\pi^+}(y) = 2f_{p\pi^0}(y)$ , with [10,17]

$$f_{p\pi^0}(y) = \frac{g^2}{16\pi^2} \frac{1}{y^2(1-y)} \int_0^\infty dk_\perp^2 |G_\pi(y, k_\perp^2)|^2 \frac{m_N^2(1-y)^2 + k_\perp^2}{[m_N^2 - M_{N\pi}^2(y, k_\perp^2)]^2}, \quad (7)$$

in which  $M_{N\pi}^2(y, k_\perp^2)$  is the invariant mass squared of the intermediate Fock state

$$M_{N\pi}^2(y, k_\perp^2) = \frac{m_N^2 + k_\perp^2}{y} + \frac{m_\pi^2 + k_\perp^2}{1-y}. \quad (8)$$

We use an exponential form for the cutoff

$$G_\pi(y, k_\perp^2) = \exp\left(\frac{m_N^2 - M_{N\pi}^2(y, k_\perp^2)}{2\Lambda^2}\right) \quad (9)$$

which insures that the identity (6) is satisfied [10,18]. We use Holtmann's parametrization [17] of the bare nucleon symmetric sea ( $\bar{Q}_{\text{bare}} = u_{\text{sea}} = \bar{u}_{\text{sea}} = d_{\text{sea}} = \bar{d}_{\text{sea}}$ )

$$x\bar{Q}_{\text{bare}}(x) = 0.11(1-x)^{15.8}. \quad (10)$$




Figure 1. Comparison of our pion cloud model with data [3] for  $\bar{d}(x) - \bar{u}(x)$ . The cutoff constant  $\Lambda = 0.83$  GeV, for which  $D = \int_0^1 (\bar{d}(x) - \bar{u}(x)) dx = 0.100$ .

Since gluon splitting is the origin of this sea, we also use a harder distribution for the bare sea quarks, of the form found in a recent determination of the gluon distribution [19]

$$x\bar{Q}'_{\text{bare}}(x) = 0.0124x^{-0.36}(1-x)^{3.8}. \quad (11)$$

For the pion valence quarks  $q_v$  and sea quarks  $q_{\text{sea}}$  we use [20]

$$xq_v(x) = 0.99x^{0.61}(1-x)^{1.02}, \quad xq_{\text{sea}}(x) = 0.2(1-x)^{5.0}. \quad (12)$$

The  $\pi$ -nucleon coupling constant is taken as  $\frac{g_\pi^2}{4\pi} = 13.6$ . The value of  $\Lambda = 0.83$  GeV is chosen to reproduce the integrated asymmetry  $D = 0.100$  [3].

### 3. Discussion

The results of our calculations for  $\bar{d}(x) - \bar{u}(x)$  are shown in Fig. 1, and those for the ratio  $\bar{d}(x)/\bar{u}(x)$  are shown in Fig. 2. In Fig. 1 the flavor asymmetry is caused entirely by the  $\pi^+$ , and our result is not affected by the different forms we have chosen for the bare sea FS contribution. In Fig. 2 the solid curve shown is for the perturbative sea quark distribution of (10). The dashed curve is for the gluonic form of (11). It is clear from Fig. 2 that a better description of the present data is provided by using a harder distribution for the symmetric sea. Of course other FS contributions could produce a similar improvement in agreement between theory and experiment. We have recently examined the role of the  $\omega$  in the meson cloud of the proton and find this to be the case [21]. The  $\sigma$  meson would also tend to suppress  $\bar{d}/\bar{u}$ . A complete calculation must include all the components of the meson cloud.

Forthcoming analyses of new E866 data [22] and proposed experiments [23] will further test our models of both perturbative and non-perturbative contributions to the flavor-symmetric and flavor-asymmetric components of the proton sea.

This work has been supported in part by the U.S. Department of Energy and the Research in Undergraduate Institutions program of the U.S. National Science Foundation (Grant No. 9722023). We thank the members of the E866 collaboration for helpful discussions of their data and analyses. One of us (EMH) thanks J. Speth for a useful discussion and for his hospitality at the KFZ in Juelich.




Figure 2. Comparison of our pion cloud model with data [3] for  $\bar{d}(x)/\bar{u}(x)$ . The solid curve is for the perturbative sea quark distribution of (10). The dashed curve is for the gluonic form of (11).

## REFERENCES

1. NMC Collaboration, P. Amaudruz et al., Phys. Rev. Lett. **66** (1991) 2712.
2. NA51 Collaboration, A. Baldit et al., Phys. Lett. **B332** (1994) 244.
3. E866 Collaboration, E.A. Hawker et al., Phys. Rev. Lett. **80** (1998) 3715; J.C. Peng et al., Phys. Rev. **D58** (1998) 092004.
4. HERMES Collaboration, K. Ackerstaff et al., Phys. Rev. Lett. **81** (1998) 5519.
5. H.L. Lai et al., Phys. Rev. **D55** (1997) 1280; A.D. Martin, R.G. Roberts and W.J. Stirling, Phys. Lett. **B387** (1996) 419.
6. A.W. Thomas, Phys. Lett. **B126** (1983) 97.
7. E.M. Henley and G.A. Miller, Phys. Lett. **B251** (1990) 453.
8. R.D. Field and R.P. Feynman, Phys. Rev. **D15** (1977) 2590.
9. W. Melnitchouk, J. Speth and A.W. Thomas, Phys. Rev. **D59** (1998) 014033.
10. J. Speth and A.W. Thomas, *Advances in Nuclear Physics*, Vol. 24, ed. J.W. Negele and E.W. Vogt (Plenum Press, NY) (1998) 83.
11. S. Kumano, Phys. Rep. **303** (1998) 183.
12. S. Kumano, Phys. Rev. **D43** (1991) 3067.
13. A. Signal, A.W. Schreiber and A.W. Thomas, Mod. Phys. Lett. **A6** (1991) 271.
14. W. Koepf, L.L. Frankfurt and M. Strikman, Phys. Rev. **D53** (1996) 2586.
15. N.N. Nikolaev, W. Schäfer, A. Szczurek and J. Speth, Phys. Rev. **D60** (1999) 014004.
16. M. Alberg, T. Falter and E.M. Henley, Nucl. Phys. **A644** (1998) 93.
17. H. Holtmann, (Mesonen im Nukleon und ihre Auswirkungen in elastischer und tiefinelastischer Streuung), Forschungszentrum Juelich, Dissertation, U. of Bonn (1995).
18. A. Szczurek, M. Ericson, H. Holtmann and J. Speth, Nucl. Phys. **A596** (1996) 397.
19. A. Vogt, hep-ph/9807369.
20. P.J. Sutton et al., Phys. Rev. **D45** (1992) 2349.
21. M. Alberg, E.M. Henley and G.A. Miller, hep-ph/9907417.
22. R. Towell, "Measurement of the Antiquark Flavor Asymmetry in the Nucleon Sea", Dissertation, University of Texas, Austin (1999).
23. P906 Collaboration, Isenhower et al., FNAL Proposal (1999).